

Akut Kalp Yetmezliđi ve Yaklaşım

Dr. Behçet Al

Chair of Emergency Medicine Department of Medicine School, Istanbul Medeniyet University/TURKEY,

Director of Emergency Department of Istanbul Göztepe Prof. Dr. Süleyman Yalçın City Hospital

Champion of ATLS and PHTLS Director Chapter of TURKEY,

AHA Instructor,

Secretary of General of Emergency Medicine Physician Association of Turkey (EPAT)

Board Member of The Muş Education and Development Foundation and Chairman of the Education Commission

Chair of Advanced Academy of Sciences Center

E-mail: behcetal@gmail.com


Mobile: +90 533 810 76 90

Office: +90 [0216 606 52 00](tel:02166065200)


Definition

- Content


- Rate and hospitalisation is increasing



Classification and Trajectories of HF Based on LVEF



Clinical assessment of the hemodynamic profile for severity



Goal of Treatment




Increase:
Quality of life




Decrease:
Symptoms,
readmission, death,
hospital stay





Treatment of HFpEF





RB indicates angiotensin receptor 2 blocker; ARNi, angiotensin receptor-neprilysin inhibitor; HF, heart failure; HFpEF, heart failure with preserved ejection fraction; LVEF, left ventricular ejection fraction; MRA, mineralocorticoid receptor antagonist; and SGLT2i, sodium-glucose cotransporter-2 inhibitor. *Greater benefit in patients with LVEF closer to 50%.

INITIAL TRIAGE

- The cornerstone is still the history and PE
- Recognition the requirement of revascularization

Diagnostic Algorithm for Patients With Suspected HF






- Address reversible factors, establish volume status and advanced medical approach

No improvement

- Try NIMV if vailable (expert and experiences)



Common Factors Precipitating HF Hospitalization With Acute Decompensated HF

ACS

Uncontrolled hypertension

AF and other arrhythmias

Additional cardiac disease (eg, endocarditis)

Acute infections (eg, pneumonia, urinary tract)

Nonadherence with medication regimen or dietary intake


Anemia

Hyper- or hypothyroidism

Medications that increase sodium retention (eg, NSAID)


Medications with negative inotropic effect (eg, verapamil)

Patients hospitalized with acute DHF



Arteriovasodilator (Na nitroglycerin, nitroprussides)

- Tachyphylaxis,
- Headache,
- 20% resistance (even high dose)
- Sudden hypotension
- Thiocyanate and cyanide toxicity,



Tests

- CBC,
- LFT
- KFT
- Eelectrolytes
- Troid FT
- Glukoz
- Lipid profile
- NT-pBNP (supportive)
- ABG
- ECG
- Bedside ECO
- Bedisde PA X-Ray

Cardiogenic shock ?

Suggested Shock Clinical Criteria

SBP <90 mm Hg for >30 min:
a. Or mean BP <60 mm Hg for >30 min
b. Or requirement of vasopressors to maintain systolic BP $\geq$ 90 mm Hg or mean BP $\geq$ 60 mm Hg
Hypoperfusion defined by:
c. Decreased mentation
d. Cold extremities, livedo reticularis
e. Urine output <30 mL/h
f. Lactate >2 mmol/L

Suggested Shock Hemodynamic Criteria

1. SBP <90 mm Hg or mean BP <60 mm Hg
2. Cardiac index <2.2 L/min/m ²
3. Pulmonary capillary wedge pressure >15 mm Hg
4. Other hemodynamic considerations <ul style="list-style-type: none">a. Cardiac power output $[(CO \times MAP)/451] < 0.6$ Wb. Shock index (HR/systolic BP) >1.0c. RV shock<ul style="list-style-type: none">i. Pulmonary artery pulse index $[(PASP-PADP)/CVP] < 1.0$ii. CVP >15 mm Hgiii. CVP-PCW >0.6

IV inotrops

Increased cardiac output and improved hemodynamics situation as needed

No measurements not improvements

- Fluid intake and output,
- Vital signs,
- Daily body weight
- Clinical signs and symptoms of congestion and
- Hypoperfusion.
- Daily tests (electrolytes, urea nitrogen, and creatinine).

Good bye?



- 33% of eligible pts with HFrEF are not prescribed ACEi-ARB-ARNi, beta blockers, and MRA therapy,
- Persistent congestion at discharge reported as 25% to 50%.

New recommendations are provided for selected pts

- Iron deficiency,
- Anemia,
- Hypertension,
- Sleep disorders,
- Type 2 diabetes,
- Atrial fibrillation,
- Coronary artery disease,
- Malignancy.

Discontinuation (beta blockers ACEi-ARB)



A higher risk for
symptoms, readmission,
rehospitalization mortality,



HOME MESSAGES

- Prevention is the shortest way
- History and PE crucial
- SGLT2i (class 2a) for HFrEF
- ARNi, ACEi, ARB, MRA, beta blockers are class 2b in this population.
- Amyloid heart disease has new recommendations for treatment
- Increased filling pressures is important for the diagnosis of HF (LVEF >40%)
- Refer patients with advanced HF to a specialized team.
- Stage A and Stage B are both **at risk**
- Figure out and treat precipitating factors
- Life style should support quality
- Don't forget to write appropriate prescription